


Provided for non-commercial research and education use.  
Not for reproduction, distribution or commercial use.



(This is a sample cover image for this issue. The actual cover is not yet available at this time.)

This article appeared in a journal published by Elsevier. The attached copy is furnished to the author for internal non-commercial research and education use, including for instruction at the author's institution and sharing with colleagues.

Other uses, including reproduction and distribution, or selling or licensing copies, or posting to personal, institutional or third party websites are prohibited.

In most cases authors are permitted to post their version of the article (e.g. in Word or Tex form) to their personal website or institutional repository. Authors requiring further information regarding Elsevier's archiving and manuscript policies are encouraged to visit:

<http://www.elsevier.com/authorsrights>



Contents lists available at ScienceDirect

Toxicon

journal homepage: [www.elsevier.com/locate/toxicon](http://www.elsevier.com/locate/toxicon)

Short communication

## Analysis of venom sac constituents from the solitary, aculeate wasp *Cerceris rybyensis*



Sachin Kote<sup>a,1</sup>, Jakub Faktor<sup>b,1</sup>, Irena Dapic<sup>a</sup>, Marcos Yebenes Mayordomo<sup>a,e</sup>,  
Mikołaj Kocikowski<sup>a,e</sup>, Alexander Kagansky<sup>c</sup>, David Goodlett<sup>a,d,h</sup>, Borivoj Vojtesek<sup>b</sup>,  
Ted Hupp<sup>a,b,e</sup>, David Wilcockson<sup>f</sup>, Ross Piper<sup>g,\*</sup>

<sup>a</sup> University of Gdansk, International Centre for Cancer Vaccine Science, ul. Wita Stwosza 63, 80-308 Gdansk, Poland

<sup>b</sup> Regional Centre for Applied Molecular Oncology, Masaryk Memorial Cancer Institute, 656 53 Brno, Czech Republic

<sup>c</sup> Centre for Genomic and Regenerative Medicine, School of Biomedicine, Far Eastern Federal University, Vladivostok, Russian Federation

<sup>d</sup> Department of Pharmaceutical Sciences, University of Maryland, Baltimore, MD, USA

<sup>e</sup> University of Edinburgh, Institute of Genetics and Molecular Medicine, Edinburgh, Scotland, EH4 2XR, UK

<sup>f</sup> IBERS, Aberystwyth University, Aberystwyth, SY23 3FL, UK

<sup>g</sup> School of Biology, University of Leeds, Leeds, LS2 9JT, UK

<sup>h</sup> Department of Microbial Pathogenesis, University of Maryland, Baltimore, MD, USA

## ARTICLE INFO

## Keywords:

Venom sac  
Mass spectrometry  
Proteomics  
Cerceris  
Hymenoptera

## ABSTRACT

Solitary aculeate wasps are abundant and diverse hymenopteran insects that disable prey using venom. The venom may possess neuromodulation, immunomodulatory, metabolic-modulatory and antimicrobial functions. Venom analysis of transcriptomes and proteomes has been previously performed in social and parasitoid wasp species. We develop methodologies including mass spectrometry-based shotgun proteomics to analyse the protein constituents from venom sacs of the solitary aculeate wasp *Cerceris rybyensis*. The venom sac constituents of *C. rybyensis* are discussed with respect to other wasp species.

### 1. Introduction

A huge number of insect species use venom to capture prey, deter predators and micro-organisms, or facilitate parasitism or extra-oral digestion (Walker et al., 2018). The bewildering diversity of insects and the multiple evolutionary origins of insect venoms represents an enormous and mostly untapped source of potentially useful therapeutic compounds, including peptides, proteins, alkaloids, and others (Walker et al., 2018). However, compared with vertebrate venoms, very little is known about those produced by insects. These venoms have been found to contain biogenic amines, peptides and proteins and are primarily for defence (Banks and Shipolini, 1986; Nakajima, 1986). These act as toxins, neuromodulators, immunomodulators, metabolic-modulators and antimicrobial agents (Konno et al., 2016).

A potentially rich hunting ground for interesting or novel venoms are the solitary aculeate wasps, of which there are around 20,000 known species (Konno et al., 2016). The ecology of these wasps is fascinating, so much so that they inspired many well-known biologists, such as Jean-Henri Fabre and Nikolaas Tinbergen (Fabre, 1916;

Tinbergen, 1932). The prey is used to provision a nest, which vary in their complexity and are stored for different durations to serve as nutrition for the developing larvae. The few studies of solitary wasp venom indicate that in addition to neurotoxins they also contain antimicrobial and cytolytic peptides, bradykinin-related peptides and neuropeptides (Konno et al., 2016). To date, the study of insect venom has been hindered by the difficulty in collecting a sufficient quantity of the venom for analysis (Konno et al., 2016). In this study we exploit state of the art shotgun mass spectrometry-based shotgun proteomic analysis to determine to what depth the venom sac proteome can be defined from the solitary wasp, *C. rybyensis*. This wasp has a palearctic distribution and is a specialist predator of solitary bees (various genera). We selected this species on the basis that its venom is known to quickly paralyse its prey, which are subsequently deposited in the warm, humid, underground nest. Adult female *C. rybyensis* wasps (Fig. 1A) were collected from a site in central England, UK. The nesting sites of the wasps were located and they were caught in flight using a net when returning to their nests with prey. The wasps were kept individually in collecting tubes and stored in a cool box. Later that day,

\* Corresponding author.

E-mail address: [ross\\_piper@yahoo.com](mailto:ross_piper@yahoo.com) (R. Piper).


<sup>1</sup> joint first author.

<https://doi.org/10.1016/j.toxicon.2019.07.012>

Received 9 May 2019; Received in revised form 19 July 2019; Accepted 23 July 2019

Available online 25 July 2019

0041-0101/ © 2019 Elsevier Ltd. All rights reserved.



**Fig. 1.** A) *Cercheris rybyensis* female (Crabronidae: Philanthinae); B) SDS-gel stained with Coomassie Blue to qualitatively estimate the amount of recovered protein in each sample pool; C) A tandem mass spectrum (MS/MS) of high-confidence VRPLPIGIPFDR peptide from trehalose-6-phosphate synthase detected in *Cercheris rybyensis* venom sacs. The figure annotates the most intensive fragment ions corresponding to the peptide and peptide coverage of trehalose-6-phosphate synthase by high confidence tryptic peptides (highlighted in green) identified in the shotgun proteomic screen of *Cercheris rybyensis* venom sacs. (For interpretation of the references to colour in this figure legend, the reader is referred to the Web version of this article.)


the wasps were anaesthetised by placing them in a  $-20\text{ }^{\circ}\text{C}$  freezer for 5 min. The venom sac was removed from each wasp under a dissecting microscope by gently pulling the sting with fine forceps while the body was held with a second pair of forceps. The extracted venom sacs were pooled into three sets of seven sacs and immediately snap frozen in microfuge tubes before storing at  $-80\text{ }^{\circ}\text{C}$ . The recovery from the sacs can be seen in Fig. 1B. The amount of protein recovered from each pool was between 280 and 420  $\mu\text{g}$ , which after extrapolation, equated to approximately 40–60  $\mu\text{g}$  of protein per sac. Approximately 25–100  $\mu\text{g}$  of protein is required for a typical shotgun proteomic experiment indicating that, in future, single animal (1 sac) experiments would be tolerated. Shotgun proteomic analysis of FASP-processed tryptic digests was performed with Eksigent Eksport nanoLC 400 (SCIEX, CA, USA) liquid chromatograph coupled to quadrupole-time-of-flight mass spectrometer TripleTOF 5600+ (SCIEX, Canada) as reported previously (Way et al., 2016). The LC-MS profile of intact crude venom extracts (Supplementary Fig. 1), their annotation and deconvolution was performed using ProteinPilot 4.5 (SCIEX, Canada). Mass spectrometry (MS) and tandem mass (MS/MS) data were searched using the redundant Uniprot + swissprot database (02.2018, 19,479 entries) restricted to Hymenoptera.

An example tandem mass spectrum of a peptide from a key protein we identified is shown in Fig. 1C. This tandem mass spectrum was shown since it contains a novel protein of importance involved in trehalose production (See below). A summary of the total proteins identified using two distinct software packages (ProteinPilot and MaxQuant) that match tandem mass spectra to sequences in a database is shown in Fig. 2A (data available on request). As expected ProteinPilot identified targets not observed with MaxQuant leading us to use the best results, sum of both informatics tools to maximally cover protein discovery. This information was the source of the pathway

characterisation described below (Fig. 2B).

Molecular chaperone proteins predominate in the venom sac, including; proteins HSP70 isoforms, HSP90, RuvBL2 (a subunit of the HSP90 containing R2TP chaperone), HSP10, BiP, HSP60, and stress-induced phospho protein-1 (Fig. 2B). Chaperones are generally involved in protein folding, protein assembly, and functional activation of protein pathways. The presence of molecular chaperones in insect venom sacs has been observed previously (Zhu et al., 2010; Colinet et al., 2014; Moreau and Asgari, 2015). Related to the maintenance of conformational integrity of proteins in the venom sac by chaperone is trehalose. Notably, our tandem mass spectra detected the enzyme trehalose-6 phosphate synthetase (Fig. 1C), which catalyses the first step in trehalose synthesis via glucose-6-phosphate. In insects, trehalose is the main storage sugar used to generate glucose for glycolysis and it is gaining interest in insect physiology as it regulates energy metabolism and glucose generation via trehalose catabolism (Nation, 2008; Shukla et al., 2015). Trehalose is known to be present in venom glands (Rivers and Denlinger, 1994a; Parkinson, 2003; Daneels et al., 2010). It has been hypothesised that venom sac trehalase converts the abundant host trehalose sugars into glucose (Rivers and Denlinger, 1994a; Parkinson, 2003), although more recent data do not support this hypothesis (Mrinalini et al., 2014). It is possible that trehalose serves as a protein stabiliser (Kaushik and Bhat, 2003; Liu et al., 2013), preserving the conformational integrity of proteins and peptides synthesised in the venom gland.

Presumably in nature, various adverse situations that protein molecules face in the venom, both within the cell and outside, can lead to protein aggregation and inactivation. This is especially true in species such as *C. rybyensis* that require high temperatures for foraging (Willmer, 1985). The use of trehalose as a natural biosynthetic agent capable of stabilizing protein molecules and helping them retain their



**Fig. 2.** A) Overview of WASP (*Cerceris rybyensis*) proteomics data analysed by MaxQuant and ProteinPilot. Quantitative Venn diagram shows, unique and common proteins between the two types of software. All the proteins were identified by shotgun proteomics approach and the dark green circle (134 proteins) indicates which are commonly identified by MaxQuant and ProteinPilot; B) *Cerceris rybyensis* venom proteins and the biochemical pathways they are involved in. (For interpretation of the references to colour in this figure legend, the reader is referred to the Web version of this article.)

functionally active forms under such conditions presumably has an adaptive role; if not a dual role as an energy source. For example, Ag-TreT1-silenced mosquitoes exhibit shorter survival under desiccation or elevated temperature (Liu et al., 2013). This latter work by Liu et al. is the first biological evidence that thermal stability can be maintained by trehalose in an animal and we would suggest this function might be important in preserving the bio-activity of venom components.

In conclusion, we report for the first time the venom sac protein constituents of the solitary aculeate wasp *C. rybyensis*. We define the yield of protein that can be recovered per gland of a solitary wasp species and the number of recovered proteins that can be detected by shotgun proteomics using two different types of discovery software. We also performed a detailed analysis of the dominant protein pathways that can be detected from these small samples. Apart from the common pathways observed in other wasp species, we also focus on the chitin biosynthetic pathway, molecular chaperone protein folding pathways, and trehalose biosynthetic pathways as dominating proteomes. We propose a novel function for trehalose production which has previously been suggested to provide a reservoir of glucose for metabolic demands. Emerging evidence suggests that trehalose can also function as an exceptional chemical chaperone and maintain the bioactive conformation of proteins. This is presumably critical for robust venom responses in host-prey interactions of *C. rybyensis*.

**Ethical statement**

Solitary wasps were the subject of this study, which are not

currently covered by legislation on the protection of animals used for scientific purposes.

**Declaration of interests**

- The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.
- The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

**Acknowledgements**

This work was supported by Ministry of Education Youth and Sports (MEYS) – NPS I - LO1413, Ministry of Health Czech Republic (MH CZ) - DRO (MMCI, 00209805). MYM and MK are part of the *Bio2Bio* study group aimed at classifying natural medicines within the PhD Doctoral Training Centre funded by the International Centre for Cancer Vaccine Science project carried out within the International Research Agendas programme of the Foundation for Polish Science co-financed by the European Union under the European Regional Development Fund.

**Appendix A. Supplementary data**

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.toxicol.2019.07.012>.

## References

- Banks, B.E.C., Shipolini, R.A., 1986. Chemistry and pharmacology of honey-bee venom. In: Piek, T. (Ed.), *Venoms of the Hymenoptera: Biochemical, Pharmacological and Behavioural Aspects*. Academic Press, London, UK, pp. 330–416.
- Colinet, D., et al., 2014. Identification of the main venom protein components of *Aphidius ervi*, a parasitoid wasp of the aphid model *Acyrtosiphon pisum*. *BMC Genomics* 15, 342.
- Daneels, E.L., et al., 2010. Venom proteins of the parasitoid wasp *nasonia vitripennis*: recent discovery of an untapped pharmacopee. *Toxins* 2, 494–516. <https://doi.org/10.3390/toxins2040494>.
- Fabre, J.-H., 1916. Translated by Alexander Teixeira de Mattos. *The hunting wasps*. Hodder and Stoughton, London.
- Kaushik, J.K., Bhat, R., 2003. Why is trehalose an exceptional protein stabilizer? An analysis of the thermal stability of proteins in the presence of the compatible osmolyte trehalose. *J. Biol. Chem.* 278 (29), 26458–26465.
- Konno, K., et al., 2016. Peptide toxins in solitary wasp venoms. *Toxins* 8 (4), 114.
- Liu, K., et al., 2013 Oct 22. Impact of trehalose transporter knockdown on *Anopheles gambiae* stress adaptation and susceptibility to *Plasmodium falciparum* infection. *Proc. Natl. Acad. Sci. U.S.A.* 110 (43), 17504–17509. <https://doi.org/10.1073/pnas.1316709110>. Epub 2013 Oct 7.
- Moreau, S.J.M., Asgari, S., 2015. Venom proteins from parasitoid wasps and their biological functions. *Toxins* 7 (7), 2385–2412.
- Mrinalini, et al., 2014. Parasitoid venom induces metabolic cascades in fly hosts. *Metabolomics* 11, 350–366.
- Nakajima, T., 1986. Pharmacological biochemistry of vespid venoms. In: Piek, T. (Ed.), *Venoms of the Hymenoptera: Biochemical, Pharmacological and Behavioural Aspects*. Academic Press, London, UK, pp. 309–327.
- Nation, J.L., 2008. *Insect Physiology and Biochemistry*. CRC Press, Boca Raton, Florida, USA.
- Parkinson, N.M., 2003. cDNAs encoding large venom proteins from the parasitoid wasp *Pimpla hypochondriaca* identified by random sequence analysis. *Comp. Biochem. Physiol.* 134, 513–520.
- Rivers, D.B., Denlinger, D.L., 1994. Developmental fate of the flesh fly, *Sarcophaga bullata*, envenomated by the pupal ectoparasitoid, *Nasonia vitripennis*. *J. Insect Physiol.* 40, 121–127.
- Shukla, E., et al., 2015. Insect trehalase: physiological significance and potential applications. *Glycobiology* 25 (4), 357–367.
- Tinbergen, N., 1932. Über die Orientierung von *Philanthus triangulum* Fabr. *Z. für Vgl. Politikwiss. (ZfVP)* 16, 305–334.
- Walker, A.A., et al., 2018. *Entomo-venomics: the Evolution, Biology and Biochemistry of Insect Venoms*, *Toxicon*. <https://doi.org/10.1016/j.toxicon.2018.09.004>.
- Way, L., et al., 2016. Rearrangement of mitochondrial pyruvate dehydrogenase subunit dihydrolipoamide dehydrogenase protein–protein interactions by the MDM2 ligand nutlin-3. *Proteomics* 16 (17), 2327–2344.
- Willmer, P.G., 1985. Size effects on the hygrothermal balance and foraging patterns of a sphecid wasp, *Cerceris arenaria*. *Ecol. Entomol.* 10, 469479.
- Zhu, J.-Y., et al., 2010. Proteomic analysis of the venom from the endoparasitoid wasp *Pteromalus puparum* (Hymenoptera: pteromalidae). *Arch. Insect Biochem. Physiol.* 75 (1), 28–44.